

# An Approach to Compute the Multi-Objective Programming Problem Under Triangular Fuzzy Numbers

**Rasha Jalal Mitlif<sup>a\*</sup>**

<sup>a</sup>Branch of Mathematics and Computer Applications, Department of Applied Sciences, University of Technology, Baghdad, Iraq.

Email: [rasha.j.mitlif@uotechnology.edu.iq](mailto:rasha.j.mitlif@uotechnology.edu.iq).

## ARTICLE INFO

### Article history:

Received: 07 /08/2022

Revised form: 20 /09/2022

Accepted : 22 /09/2022

Available online: 24 /09/2022

### Keywords:

Multi- objective function (MOF),  
 Triangular fuzzy number (TrFN),  
 Fractional programming problems  
 (FrPP), Linear Programming  
 Problem (LnPP) , Simplex method  
 (SM).

## ABSTRACT

The most important one advantages of the operation research (OR) theory is displayed of this paper for tackling multi-objective function programming problems (MOFPP). In this study, we present an approach for multi-objective fractional programming problems (MOFrPP) using fuzzy numbers (FN) coefficients in the objective functions (OF). The objective functions' (OF) parameters are all considered to be fuzzy triangular fuzzy numbers (FTrFN). This problem's multi-objective function (MOF) case was resolved by transforming the MOF to a linear programming problem (LnPP) that could be resolved using the simplex method (SM). The problem is being solved using algorithms. Then the value is compared with a result, which obtained by simplex method (SM). These techniques are illustrated with numerical examples (NE). This study shows the reliability of our methods and their utility to this specific set of multi-objective functions (MOF).

MSC..

<https://doi.org/10.29304/jqcm.2022.14.3.1034>

## 1. Introduction

The method of multi-objective function (MOF) is useful for analyzing the best trade-off solutions that balance various criteria. It is a little challenging to maximize two or more goals at once, and the challenge increases if the goals are incompatible. It was intended to analyze the potential for coming up with a compromise solution that achieves all the goals at once. Several techniques have been introduced for solving linear problems (LP). Some of them are extensions of the fractional linear problems (FrLP) and others are based on different idea. Mediya B. Mrakhan et al. [1] a new approach to the scaling-based multi-objective linear programming problem (MOLrPP). Ronak M. Abdullah and Snoor O.Abdalla [2] studied solutions of MOLrPP by Advanced Transformation Technique. Samsun Nahar and Md. Abdul Alim [3] presented a new technique to solve a MOLnFrPP by using new geometric averaging method and used some other techniques such as arithmetic averaging, geometric averaging. Farhana Akond Pramy and Md. Ainul Islam [4] the multi-objective linear fractional programming problem (MOLnFrP) problem was given a single efficient solution, and finds the effective solutions of (MOLnFrPP). S. F. Tantawy [5]

\*Corresponding author Rasha Jalal Mitlif

Email addresses: [rasha.j.mitlif@uotechnology.edu.iq](mailto:rasha.j.mitlif@uotechnology.edu.iq).

Communicated by 'sub etitor'

Described a feasible direction be used to locate all effective extreme points for MOLnFrPP when all denominators are equal. Moslem Ganji and Mansour Sara [6] showed the relationship between its Robust Counterpart (RC) formulations and a Multi Objective Linear Fractional Programming (MOLrFrP) challenge with uncertain data in the objective function (OF). R. Suganya and U. Afrin [7] provided a new statistical averaging method (New Quadratic mean) for MOLrPP. [8-16] A number of fuzzy numbers have been used to solve MOFrPP by many researchers.

In order to improve this work, we established a MOLrFrPP, offered an algorithm to resolve the fractional programming problem for the multi-objective function, and proposed a creative approach to resolve the issue in order to produce the best optimal solution. By resolving a numerical example, the computer application of our approach has also been explained. Finally, we presented the findings and their comparability. Some basic definitions have been proposed, which will be important in this paper.

### Definition 2.1: Fuzzy Number (FN) [17]

If the membership function is satisfied, the FN  $\tilde{E}$  is fuzzy

(i) A fuzzy set of the discourse  $G$  is convex

(ii)  $\tilde{E}$  is normal if  $\exists g_i \in G, M_{\tilde{E}}(g_i) = 1$

(iii) and  $M_{\tilde{E}}(b)$  is piecewise incessant.

### Definition 2.2: Triangular Fuzzy Number (TrFN) [18]

A TrFN is defined as  $\tilde{J} = (\tau, \eta, \varepsilon)$  for  $\tau, \eta, \varepsilon \in \mathbb{R}$  and its membership is given by

$$M_{\tilde{J}}\tilde{h} = \begin{cases} \frac{\tilde{h}-\tau}{\eta-\tau} & \tau \leq \tilde{g} \leq \eta \\ \frac{\varepsilon-\tilde{h}}{\varepsilon-\eta} & \eta \leq \tilde{g} \leq \varepsilon \\ 0 & \text{otherwise} \end{cases}$$

### 2. Linear Programming Problem (LnPP) [19]

A LnPP of the form

Maximize  $c^t g$

Subject to

$Og = \bar{b}$

And

$g \geq 0$

Where  $c \in \hat{Y}^n$ ,  $\bar{b} \in \hat{Y}^n$  and  $O \in \hat{Y}^m \cup^n$ .

### 3. Linear Fractional Programming Problem (LnFrPP) [20]

The general linear fractional programming problem is defined as follows:

$$\text{Maximize } \check{Z} = \frac{T(q)}{3(q)}$$

$T(q) = c^t q + \alpha$  ,  $3(q) = d^t q + \beta$  are valued and continuous functions on  $U$  and  $d^t q + \beta \neq 0$  for all  $q \in U$  and  $U = \{q: Oq = \bar{b}, q \geq 0\}$ ,  $q, c^t, d^t \in R^n$  ,  $\alpha, \beta \in \hat{Y}$   $\bar{b} \in R^m$  ,  $O \in \hat{Y}^{m \times n}$  .

#### 4. Multi-Objective function Technique (MOFT) [21]

The mathematical Multi-Objective function Technique (MOFT) [21] is explained as:

$$\text{Optimal Max } \check{Z} = [\text{Max } Z_1 , \text{Max } Z_2 , \text{Max } Z_3, \dots]$$

Subject to:

$$Oq = \bar{b}, q \geq 0$$

#### 5. Complementary Multi-Objective Fractional Programming Formula (CMOFrPP)

The Complementary MOLnFrPP method have been used by the scientists [22-23] to solve the multi-objective (MO) optimization problems.

#### 6. Ranking function (RF)

In a fuzzy numbers the ranking function is essential in the process for determining decisions. Ranking of FN is required to find the largest and smallest fuzzy numbers. The article [24] have proposed ranking function methods as the follows:

$$H(\tilde{V}) = \frac{\xi + 14 * t + b}{16} \dots (1)$$

#### 7. The New Algorithm for Solving multi-objective function programming problems (MOFPP)

An algorithm to explain MOFPP by SM which can be summarized as follows:

Step 1: We suggest two techniques for process of the converting fuzzy multi-objective fractional functions to linear fractional functions programming problems (FMOLrFrPP) into LPP.

Step2: At first consider the OF *Maximize*  $Z = \frac{T(q)}{3(q)}$  is FMOLrFPP.

Step3: FMOLrFPP is changed to the LrFPP by two techniques: the first technique is complementary fractional and the second technique is ranking function.

Step4: dividing the first fractional objective function (FrOF) into two linear functions (LnF), the first function serving as the numerator function (NF) and the second as the denominator function (DF). Maximum  $T(q)$  for the numerator and minimize  $3(q)$  for the denominator functions are used to determine the objective function's value.

Step 5: An LrP problem is written as  $Max Z^*$  subject to the original problem's restrictions, where  $Max Z^*$  is the result of subtracting the denominator function  $\mathfrak{Z}(g)$  from the numerator function  $\mathfrak{T}(g)$ , or  $Maximize Z = \mathfrak{T}(g) - \mathfrak{Z}(g)$ .

Step 6: The second, third, and fourth objective functions all follow the same process.

Step 7: Find the optimal solution by using Simplex process of each objective function.

The algorithm is illustrated with the following example.

### 8. Numerical Example :

The following example is illustrated to show the proposed methodology for MOFFr the following MOFFr is formulated as

$$\text{Max } Z_1 = \frac{(6,7,8) \gamma_1 + (4,5,6) \gamma_2}{(4,5,6) \gamma_1 + (3,4,5) \gamma_2}$$

$$\text{Max } Z_2 = \frac{(7,8,9) \gamma_1 + (4,5,6) \gamma_2}{(6,7,8) \gamma_1 + (3,4,5) \gamma_2}$$

$$\text{Max } Z_3 = \frac{(8,9,10) \gamma_1 + (5,6,7) \gamma_2}{(5,6,7) \gamma_1 + (2,3,4) \gamma_2}$$

subject to

$$(4,5,6) \gamma_1 + (6,7,8) \gamma_2 \leq (16,17,18)$$

$$(5,6,7) \gamma_1 + ((8,9,10) \gamma_2 \leq (20,21,22)$$

$$(6,7,8) \gamma_1 + (3,4,5) \gamma_2 \leq (11,12,13)$$

$$\gamma_1, \gamma_2 \geq 0.$$

1. First technique: The above problem can be converted with all the three maximization fuzzy multi-objective fractional functions to linear fractional functions as detailed below:

$$\text{Max } Z_1 = \frac{8 \gamma_1 + 6 \gamma_2}{6 \gamma_1 + 5 \gamma_2}$$

$$\text{Max } Z_2 = \frac{9 \gamma_1 + 6 \gamma_2}{8 \gamma_1 + 5 \gamma_2}$$

$$\text{Max } Z_3 = \frac{10 \gamma_1 + 7 \gamma_2}{7 \gamma_1 + 4 \gamma_2}$$

subject to

$$6 \gamma_1 + 8 \gamma_2 \leq 18$$

$$7 \gamma_1 + 10 \gamma_2 \leq 22$$

$$8 \gamma_1 + 5 \gamma_2 \leq 13$$

$$Y_1, Y_2 \geq 0.$$

2. Now the new objective function  $Max Z^*$  is constructed as per the complementary fractional method.

$$Max Z_1^* = 2 Y_1 + Y_2$$

$$Max Z_2^* = Y_1 + Y_2$$

$$Max Z_3^* = 3 Y_1 + 3 Y_2$$

subject to

$$6 Y_1 + 8 Y_2 \leq 18$$

$$7 Y_1 + 10 Y_2 \leq 22$$

$$8 Y_1 + 5 Y_2 \leq 13$$

$$Y_1, Y_2 \geq 0.$$

We solve first linear objective function to obtain the optimal solution.

$$Max Z_1^* = 2 Y_1 + Y_2$$

subject to

$$6 Y_1 + 8 Y_2 \leq 18$$

$$7 Y_1 + 10 Y_2 \leq 22$$

$$8 Y_1 + 5 Y_2 \leq 13$$

$$Y_1, Y_2 \geq 0.$$

Solving by Simplex Method using (Win QSB) Programming, we get

$$Max Z_1^* = 3.250, Y_1 = 1.625, Y_2 = 0$$

We solve second linear objective function to obtain the optimal solution.

$$Max Z_2^* = Y_1 + Y_2$$

subject to

$$6 Y_1 + 8 Y_2 \leq 18$$

$$7 Y_1 + 10 Y_2 \leq 22$$

$$8 Y_1 + 5 Y_2 \leq 13$$

$$Y_1, Y_2 \geq 0.$$

Solving by Simplex Method using (Win QSB) Programming, we get

$$Max Z_2^* = 2.333, Y_1 = 0.444, Y_2 = 1.888$$

We solve third linear objective function to obtain the optimal solution.

$$\text{Max } Z_3^* = 3 \gamma_1 + 3 \gamma_2$$

subject to

$$6 \gamma_1 + 8 \gamma_2 \leq 18$$

$$7 \gamma_1 + 10 \gamma_2 \leq 22$$

$$8 \gamma_1 + 5 \gamma_2 \leq 13$$

$$\gamma_1, \gamma_2 \geq 0.$$

Solving by Simplex Method using (Win QSB) Programming, we get

$$\text{Max } Z_3^* = 7.000, \gamma_1 = 0.444, \gamma_2 = 1.888$$

Second technique: the ranking function of triangular fuzzy numbers  $H(\tilde{V}) = \frac{\xi+14*\zeta+b}{16}$ , the above fuzzy multi-objective fractional functions (FMOFLFPP) can be converted to the following multi-objective fractional functions (MOLFPP):

$$\text{Max } Z_1 = \frac{7 \gamma_1 + 5 \gamma_2}{5 \gamma_1 + 4 \gamma_2}$$

$$\text{Max } Z_2 = \frac{8 \gamma_1 + 5 \gamma_2}{7 \gamma_1 + 4 \gamma_2}$$

$$\text{Max } Z_3 = \frac{9 \gamma_1 + 6 \gamma_2}{6 \gamma_1 + 3 \gamma_2}$$

subject to

$$6 \gamma_1 + 8 \gamma_2 \leq 18$$

$$7 \gamma_1 + 10 \gamma_2 \leq 22$$

$$8 \gamma_1 + 5 \gamma_2 \leq 13$$

$$\gamma_1, \zeta_2 \geq 0.$$

We solve three linear programming problems to obtain the optimal solution.

$$\text{Max } Z_1^* = 2 \gamma_1 + \gamma_2$$

$$\text{Max } Z_2^* = \gamma_1 + \gamma_2$$

$$\text{Max } Z_3^* = 3 \gamma_1 + 3 \gamma_2$$

subject to

$$6 \gamma_1 + 8 \gamma_2 \leq 18$$

$$7 \gamma_1 + 10 \gamma_2 \leq 22$$

$$8 \gamma_1 + 5 \gamma_2 \leq 13$$

$$Y_1, Y_2 \geq 0.$$

We solve first linear objective function to obtain the optimal solution.

$$\text{Max } Z_1^* = 2 Y_1 + Y_2$$

subject to

$$6 Y_1 + 8 Y_2 \leq 18$$

$$7 Y_1 + 10 Y_2 \leq 22$$

$$8 Y_1 + 5 Y_2 \leq 13$$

$$Y_1, Y_2 \geq 0.$$

Solving by Simplex Method using (Win QSB) Programming, we get

$$\text{Max } Z_1^* = 3.250, Y_1 = 1.625, Y_2 = 0$$

We solve second linear objective function to obtain the optimal solution.

$$\text{Max } Z_2^* = Y_1 + Y_2$$

subject to

$$6 Y_1 + 8 Y_2 \leq 18$$

$$7 Y_1 + 10 Y_2 \leq 22$$

$$8 Y_1 + 5 Y_2 \leq 13$$

$$Y_1, Y_2 \geq 0.$$

Solving by Simplex Method using (Win QSB) Programming, we get

$$\text{Max } Z_2^* = 2.333, Y_1 = 0.444, Y_2 = 1.888$$

We solve third linear objective function to obtain the optimal solution.

$$\text{Max } Z_3^* = 3 Y_1 + 3 Y_2$$

subject to

$$6 Y_1 + 8 Y_2 \leq 18$$

$$7 Y_1 + 10 Y_2 \leq 22$$

$$8 Y_1 + 5 Y_2 \leq 13$$

$$Y_1, Y_2 \geq 0.$$

Solving by Simplex Method using (Win QSB) Programming, we get

$$\text{Max } Z_3^* = 7.000, Y_1 = 0.444, Y_2 = 1.888$$

the results of above example when solved by complementary fractional method and ranking function method almost it same result are shown below in the table(1).

**Table (1): Resulting multi-objective function programming problems (MOFPP):**

3. No	4. Method	5. The optimal solution (OS)
6. 1	7. complementary fractional method	8. $Max Z_1^* = 3.250, y_1 = 1.625, y_2 = 0$ 9. 10. $Max Z_2^* = 2.333, y_1 = 0.444, y_2 = 1.888$ 11. 12. $Max Z_3^* = 7.000, y_1 = 0.444, y_2 = 1.888$ 13.
14. 2	15. Ranking function	16. $Max Z_1^* = 3.250, y_1 = 1.625, y_2 = 0$ 17. 18. $Max Z_2^* = 2.333, y_1 = 0.444, y_2 = 1.888$ 19. 20. $Max Z_3^* = 7.000, y_1 = 0.444, y_2 = 1.888$ 21.

## 9. Conclusion

A new TrFN algorithm to convert the MOLPP into a single LPP. Additionally, in order to compare the outcomes, we applied our strategy to the example using two distinct approaches to the problem. This comparison allowed us to see that two method produced results that were exactly the same value and its computer application employing programming in the mathematical language (Win QSB).

## References

- [1] Mediya B. Mrakhan , Ayad M. Ramadan , Rzgar F. Mahmood, and Ronak M. Abdullah , A NOVEL TECHNIQUE FOR SOLVING MULTIOBJECTIVE LINEAR PROGRAMMING PROBLEMS , *Mathematical Forum* , 28(2), 2020 , 155-165.
- [2] Ronak M. Abdullah, Snoor O. Abdalla , Solution of Multi-Objective Linear Programming Problem by Advanced Transformation Technique, *Samarra Journal of Pure and Applied Science, Samarra J. Pure Appl. Sci.*, 2022; 4 (2): 118-128.
- [3] Samsun Nahar and Md. Abdul Alim , A New Geometric Average Technique to Solve Multi-Objective Linear Fractional Programming Problem and Comparison with New Arithmetic Average Technique , *IOSR Journal of Mathematics (IOSR-JM)* ,13(3), 2017,39-52.
- [4] Farhana Akond Pramy, Md. Ainul Islam , Determining Efficient Solutions of Multi-Objective Linear Fractional Programming Problems and Application , *Open Journal of Optimization*, 2017, 6, 164-175.
- [5] S. F. TANTAWY, SOLVING A SPECIAL CLASS OF MULTIPLE OBJECTIVE LINEAR FRACTIONAL PROGRAMMING PROBLEMS, the ANZIAM journal, 56, 2014, 91–103.
- [6] Moslem Ganji , Mansour Saraj , Solving Multi Objective Linear Fractional Programming Problem Under Uncertainty via Robust Optimization Approach , *Journal of Informatics and Mathematical Sciences* , Vol. 11 No. 2 , 2019, 115-125.
- [7] R. Suganya and U. Afrin , SOLVING MULTI-OBJECTIVE LINEAR PROGRAMMING PROBLEMS BY STATISTICAL AVERAGING AND NEW STATISTICAL AVERAGING METHOD, *International Journal of Mechanical Engineering* , Vol. 7 No. 4 April, 2022 , 1038-1043.
- [8] S. Muruganandam and P. Ambika , Harmonic Mean Technique to Solve Multi Objective Fuzzy Linear Fractional Programming Problems, *Global Journal of Pure and Applied Mathematics* , Volume 13, Number 10, (2017), pp. 7321-7329.
- [9] Beena T Balan ,Solving Multi-Objective Fuzzy Linear Optimization Problem Using Fuzzy Programming Technique, *IOSR Journal of Mathematics (IOSR-JM)* , 2016, 18-21.
- [10] P. A. Thakre, D. S. Shelar , S. P. Thakre , Solving fuzzy linear programming problem as multi objective linear programming problem, *Journal of Engineering and Technology Research* Vol. 2(5), pp. 82-85, 2009.
- [11] Xiao-Peng Yang , Bing-Yuan Cao , Hai-Tao Lin , Multi-Objective Fully Fuzzy Linear Programming Problems with Triangular Fuzzy Numbers, 11th International Conference on Fuzzy Systems and Knowledge Discovery (FSKD), 2014, 165-171.
- [12] Nejmaddin A. Sulaiman, Rebaz B. Mustafa , Using Harmonic Mean to Solve Multi-Objective Linear Programming Problems , *American Journal of Operations Research*, 2016, 6, 25-30.
- [13] .Rasha Jalal Mitlif, A New Method for Solving Fully Fuzzy Multi-Objective Linear Programming Problems , *Iraqi Journal of Science*, 2016, Vol. 57, No.3C, pp:2307-2311.
- [14] Rasha Jalal Mitlif , Solving fuzzy fractional linear programming problems by ranking function methods , *JOURNAL OF COLLEGE OF EDUCATION*, 2016, NO.1 , 93-108.



- [15] I. H. Hussein and R. J. Mitlif, "Ranking Function to Solve a Fuzzy Multiple Objective Function", *Baghdad Science Journal*, vol. 18, no. 1, pp. 144-148 , 2021.
- [16] Purnima Pandit , Multi-objective Linear Programming Problems involving Fuzzy Parameters, *International Journal of Soft Computing and Engineering (IJSCE)*, Volume-3, Issue-2, 2013 , 99-102.
- [17] N. RAMESHAN and D. STEPHEN DINAGAR , A STUDY ON FUZZY CRITICAL PATH WITH QUADRILATERAL FUZZY NUMBERS , *Advances and Applications in Mathematical Sciences* 21(3), 2022, 1491-1508 .
- [18] Vijay C. Makwana, Vijay P. Soni, Nayan I. Patel, Manoj Sahni, Fuzzy Number – A New Hypothesis and Solution of Fuzzy Equations , *Mathematics and Statistics* 10(1): 176-186, 2022.
- [19] Nejmaddin A. Suleiman and Maher A. Nawkhass , A New Modified Simplex Method to Solve Quadratic Fractional Programming Problem and Compared it to a Traditional Simplex Method by Using Pseudoaffinity of Quadratic Fractional Functions, *Applied Mathematical Sciences*, Vol. 7, 2013, no. 76, 3749 – 3764.
- [20] El- Saeed Ammar, Mohamed Muamer, Algorithm for Solving Multi Objective Linear Fractional Programming Problem with Fuzzy Rough Coefficients, *Open Science Journal of Mathematics and Application* , 2016; 4(1): 1-8.
- [21] Chandra Sen , Duality in Solving Multi-Objective Optimization (MOO) Problems, *American Journal of Operations Research*, 2019, 9, 109-113.
- [22] T. Ibaraki, Complementary programming, *Oper. Res.* 19 (1971), 1523–1529.
- [23] Maher Nawkhass , Nejmaddin Sulaiman , Symmetric fuzzy approach to solve complementary multi-objective linear fractional programming problem , *Int. J. Nonlinear Anal. Appl.* In Press, 2022, 1–6.
- [24] Rasha Jalal Mitlif , An Efficient Algorithm for Fuzzy Linear Fractional Programming Problems via Ranking Function , *Baghdad Science Journal* , 2022, 19(1): 71-76.