

Topological Spaces

Rewayda Razaq Mohsin Ahmed M. Rajab Enas Y. Abdalla
Coll. of Math. Sci. Math-Dep. Coll. of Edu. Math-Dep.
University of Kufa, Najaf, Iraq

Abstract:

The concept of intuitionistic (double) topological spaces was introduced by Çoker 1996. The aim of this paper is to give a notion of pairwise compactness for double topological spaces and some separation axioms .

المخلص:

فكرة فضاء التبولوجي المضاعف قدم من قبل Coker في العام 1996 . الهدف من هذا البحث هو تقديم تعريف التراص الزوجي للفضاءات التبولوجية المضاعفة وبعض بديهيات الفصل.

1.Introduction

The concept of a fuzzy topology was introduced by Chang in 1968 [2] after the introduction of fuzzy sets by Zadeh in 1965. Later this concept was extended to intuitionistic fuzzy topological spaces by Çoker in [4] . In [5] Coker studied continuity, connectedness, compactness and separation axioms in intuitionistic fuzzy topological spaces. In this paper we follow the suggestion of J.G. Garcia and S.E. Rodabaugh [7] that (double fuzzy set) is a more appropriate name than (intuitionistic fuzzy set) ,and therefore adopt the term (double-set) for the intuitionistic set , and (double-topology) for the intuitionistic topology of Dogan Çoker , (this issue) we denote by **Dbl-Top** the construct (concrete texture over Set) whose objects are pairs (X, τ) where τ is a double-topology on X .In Section three we discuss making use of this relation between bitopological spaces and double- topological spaces , we generalize a notion of compactness for double- topological space in section four with some theorems about T_1 , T_2 , T_3 .

2.Preliminaries

Throughout the paper by X we denote a non-empty set . In this section we shall present various fundamental definitions and propositions. The following definition is obviously inspired by Atanassov [1].

2.1. Definition

[8] A double-set (Ds in brief) A is an object having the form $A = \langle X, A_1, A_2 \rangle$. Where A_1 and A_2 are subsets of X satisfying $A_1 \cap A_2 = \phi$. The set A_1 is called the set of members of A , while A_2 is called the set of non-members of A .

Throughout the remainder of this paper we use the simpler $A = (A_1, A_2)$ for a double-set.

2.2. Remark

Every subset A of X may obviously be regarded as a double-set having the form $A = (A, A^c)$, where $A^c = X \setminus A$ is the complement of A in X . We recall several relations and operations between DS's as follows:

2.3. Definition

[8] Let the DS's A and B on X be the form $A = (A_1, A_2)$, $B = (B_1, B_2)$, respectively. Furthermore, let $\{A_j : j \in J\}$ be an arbitrary family of DS's in X , where $A_j = (A_j^{(1)}, A_j^{(2)})$. Then

- (a) $A \subseteq B$ if and only if $A_1 \subseteq B_1$ and $A_2 \supseteq B_2$;
- (b) $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$;
- (c) $\bar{A} = (A_2, A_1)$ denotes the complement of A ;
- (d) $\bigcap A_j = (\bigcap A_j^{(1)}, \bigcup A_j^{(2)})$;
- (e) $\bigcup A_j = (\bigcup A_j^{(1)}, \bigcap A_j^{(2)})$;
- (f) $\square A = (A_1, A_1^c)$;
- (g) $\langle \rangle A = (A_2^c, A_2)$;
- (h) $\phi = (\phi, X)$ and $X = (X, \phi)$.

In this paper we require the following :

- (i) $(\)A = (A_1, \phi)$, and (ii) $(\)A = (\phi, A_2)$.

Is call the image and preimage of DS's under a function.

2.4. Definition. [3,8]

Let $x \in X$ be a fixed element in X . Then:

- (a) The DS given by $\tilde{x} = (\{x\}, \{x\}^c)$ is called a double-point (DP in brief X).
- (b) The DS $\tilde{x} = (\phi, \{x\}^c)$ is called a vanishing double-point (VDP in brief X).

2.5.Definition. [3,8]

(a) Let \tilde{x} be a DP in X and $A=(A_1,A_2)$ be a DS in X . Then $\tilde{x} \in A$ iff $x \in A_1$.

(b) Let \tilde{x} be a VDP in X and $A=(A_1,A_2)$ a DS in X . Then $\tilde{x} \in A$ iff $x \notin A_2$.

It is clear that $\tilde{x} \in A \Leftrightarrow x \subseteq A$ and that $\tilde{x} \in A \Leftrightarrow x \subseteq A$.

2.6.Definition

[10] A double-topology (DT in brief) on a set X is a family τ of DS's in X satisfying the following axioms :

T1: $\phi, X \in \tau$,

T2: $G_1 \cap G_2 \in \tau$ for any $G_1, G_2 \in \tau$,

T3: $\cup G_j \in \tau$ for any arbitrary family $\{G_j : j \in J\} \subseteq \tau$.

In this case the pair (X, τ) is called a double-topological space (DTS in brief), and any DS in τ is known as a double open set (DOS in brief). The complement \bar{A} of a DOS A in a DTS is called a double closed set (DCS in brief) in X .

2.7.Definition

[10] Let (X, τ) be an DTS and $A = (A_1,A_2)$ be a DS in X.

Then the interior and closure of A are defined by :

$$\text{int}(A) = \cup \{G : G \text{ is a DOS in } X \text{ and } G \subseteq A\},$$

$$\text{cl}(A) = \cap \{H : H \text{ is a DCS in } X \text{ and } A \subseteq H\},$$

respectively .

It is clear that $\text{cl}(A)$ is a DCS in X and $\text{int}(A)$ a DOS in X . Moreover A is a DCS in X iff $\text{cl}(A) = A$, and A is a DOS in X iff $\text{int}(A) = A$.

2.8. Example

[5] Any topological space (X, τ_0) gives rise to a DT of the form $\tau = \{A' : A \in \tau_0\}$ by identifying a subset A in X with its counterpart $A' = (A, A^c)$, as in Remark 2.2.

3- The Construction of Dbl-Top and Bitop :

We begin by recalling the following results which associates a bitopology with a double topology.

3.1. Proposition

[5] Let (X, τ) be a DTS.

- (a) $\tau_1 = \{A_1 : \exists A_2 \subseteq X \text{ with } A = (A_1, A_2) \in \tau\}$ is a topology on X .
 (b) $\tau_2^* = \{A_2 : \exists A_1 \subseteq X \text{ with } A = (A_1, A_2) \in \tau\}$ is the family of closed sets of the topology $\tau_2 = \{A_2^c : \exists A_1 \subseteq X \text{ with } A = (A_1, A_2) \in \tau\}$ is a topology on X .
 (c) Using (a) and (b) we may conclude that (X, τ_1, τ_2) is a bitopological space.

3.2. Proposition

Let (X, u, v) be a bitopological space. Then the family

$$\{(U, V^c) : U \in u, V \in v, U \subseteq V\}$$

is a double topology on X .

Proof : The condition $U \subseteq V$ ensures that $U \cap V^c = \phi$, while the given family contains ϕ because $\phi \in u, v$, and it contains X because $X \in u, v$. Finally this family is closed under finite intersections and arbitrary unions by Definition 2.3 (d,e) and the corresponding properties of the topologies u and v .

3.3. Definition

Let (X, u, v) be a bitopological space. Then we set

$$\tau_{uv} = \{(U, V^c) : U \in u, V \in v, U \subseteq V\}$$

and call this the double topology on X associated with (X, u, v) .

3.4. Proposition

If (X, u, v) is a bitopological space and τ_{uv} the corresponding DT on X , then

$$(\tau_{uv})_1 = u \text{ and } (\tau_{uv})_2 = v.$$

Proof. $U \in u$ implies $(U, \phi) \in \tau_{uv}$ since $U \subseteq X \in v, \text{ so } (U, \phi) \in \tau_{uv}$. Conversely, take $(U, B) \in \tau_{uv}$. Then $(U, B) \in \tau_{uv}$ for some $B \subseteq X$, and now $U \in u$. Hence $(\tau_{uv})_1 \subseteq u$, and the first equality is proved.

The proof of the second equality may be obtained in a similar manner, and we omit the details. Now we define double compact set and we use the link between bitopological space and double topological space to established some theorems .

4.Pairwise Double Compact Set :

4.1. Definition

By an double open cover of a subset A of a double topological space (X, τ) , we mean a collection $C = \{G_j : j \in J\}$ of double open subsets of X such that $A \subset \bigcup \{G_j : j \in J\}$ then we say that C covers A . In particular , A collection C is said to be an open cover of the space X iff $X = \bigcup \{(G_j^1, G_j^2) : j \in J\}$ of double open subsets of X .

4.2.Definition

A double-set A of DTS in (X, τ) is said to be double compact set iff every double sub cover , that is iff for every collection $\{G_j : j \in J\}$ of DOS's for which $A \subset \{G_j : j \in J\}$ for $A = (A_1, A_2)$ such that $(A_1, A_2) \subset (G_{j_1}^1, G_{j_1}^2) \cup \dots \cup (G_{j_n}^1, G_{j_n}^2)$.

4.3.Definition

Let (X, τ) be DTS and let $N \in X$. A double set N of X is said to be τ -nhd of \tilde{x} iff there exists τ -DOS , G such that $\tilde{x} \in G \subset N$, similarly N is called a τ -double nhd of $A \subset X$ iff there exists an DOS , G such that $A \subset G \subset N$.

4.4.Definition

[3] The DTS (X, τ) is called pairwise T_2 if given $x \neq y$ in X there exists $G, H \in \tau$ Satisfying $\tilde{x} \in G, \tilde{y} \in H$ and $G \subseteq \overline{H}$.

4.5. Proposition

If (X, τ) is pairwise T_2 then every double compact set is double closed set.

Proof: We shall show that $\bar{G} \in \tau$ is double open set. Let $p \in \bar{G} = (G_1, G_2)$.

Since X is T_2 then for.

Then $p \in G_2, y \notin H_2 \Rightarrow y \in H_2^c, G_2 \cap H_2^c = \phi$

\exists double open nhds of $p, y, \mu(p)$ & $N(y)$ such that $\mu(p) \cap N(y) = \phi$

Now the collection $\{\mu(p) : p \in G_2\}$ double open cover of G_2

$\because G$ is compact then $\{G_2 \subset \cup \mu(p_i)\}$.

let $M = \cup \mu(p_i), N = \cap N(y_i)$ then N is double open nhd of y_i

We claim that $M \cap N = \phi$,

$z \in \mu \Rightarrow z \in \mu(p_i) \Rightarrow z \notin N(y_i) \Rightarrow z \notin N$, thus $M \cap N = \phi$

Since $G_2 \subset M$, then $G_2 \cap N = \phi \Rightarrow N \subset G_2 \Rightarrow N \subset \bar{G}$ this shows that \bar{G} contains a nhd of each of its point and so \bar{G} is DOS otherwise G is DCS.

4.6. Proposition

Let A and B be disjoint double compact subsets of a DTS (X, τ)

Then there exists disjoint DOS's G and H such that $A \subset G$ and $B \subset H$.

Proof : First, let $x \in A$ be fixed. Since X is pairwise T_2 and $x \notin B$, for each $y \in B$

$A \subseteq (\bar{B})^c$. (clearly $x \in A_1, y \notin B_2 \Rightarrow y \in B_2^c$ for $A = (A_1, A_2), B = (B_1, B_2)$)

There exists DOS's G_y and H_y such that $x \in G_y$ and $y \in H_y$. The

collection $\{H_y : y \in B\}$ is a double open cover of B . Since B is double compact

subspace of X , there exist finitely many points y_1, y_2, \dots, y_n of B such that

$$B \subset \{H_{y_i} : i = 1, 2, \dots, n\}, (B_1, B_2) \subset \{(H_{y_i}^1, H_{y_i}^2) : i = 1, 2, \dots, n\}$$

$$\text{let } G_x = \cap \{G_{y_i} : i = 1, 2, \dots, n\} = \cap \{(G_{y_i}^1, G_{y_i}^2) : i = 1, 2, \dots, n\}$$

$H_x = \cup \{(H_{y_i}^1, H_{y_i}^2) : i = 1, 2, \dots, n\}$ then G_x, H_x are disjoint open sets such

that $x \in G_x$ and $B \subset H_x$.

now let $\tilde{x} \in A$ be arbitrary and let \tilde{G}_x and \tilde{H}_x be as constructed above, then evidently the collection $\{\tilde{G}_x : x \in A\}$ is a double open cover of A . Since A is a double compact subspace of X . There exist finitely many points, $\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_m$ such that $A \subset \bigcup\{\tilde{G}_{\tilde{x}_i} : i = 1, 2, \dots, m\}$, let $\tilde{G} = \bigcup\{\tilde{G}_{\tilde{x}_i} : i = 1, 2, \dots, m\}$ and $\tilde{H} = \bigcap\{\tilde{H}_{\tilde{x}_i} : i = 1, 2, \dots, m\}$ then \tilde{G} and \tilde{H} are disjoint double open sets such that
 $\tilde{A} \subset \tilde{G}$ and $\tilde{B} \subset \tilde{H}$.

4.7. Definition

[3] The DTS (X, τ) is called pairwise T_1 if given $x \neq y$ in X there exists $G \in \tau$ with $x \in G, y \notin G$, and there exists $H \in \tau$ with $y \in H, x \notin H$.

4.8. Definition

[6] The DTS (X, τ) is called pairwise T_3 if $\forall DCS A \in \tau, a \in \text{int } A$ in X there exists $G, H \in \tau$ satisfying $a \in G, a \notin H, A \subseteq H$ and $G \subseteq \overline{H}$.

4.9. Proposition

The DTS (X, τ) is called pairwise T_1 iff every singleton double set $\{\tilde{x}\}$ of X is DCS.

Proof : \Leftarrow Let every singleton double set $\{\tilde{x}\}$ of X be DCS to show that the space

is T_1 . Let x, y be any two disjoint double point of X , then $\{\tilde{x}\}$ is a DOS which contain y

Similarly $\{\tilde{y}\}$ is a DOS which contain x but does not contain y . Hence (X, τ) is pairwise T_1 .

\Rightarrow Let the space be pairwise T_1 and let x be any point of X , we want to show that $\{\tilde{x}\}$ is DCS, that to show $X - \{\tilde{x}\}$ is DOS. Let $\tilde{y} \in X - \{\tilde{x}\}$ then $x \neq y$ since X is pairwise T_1 .

There exist an open G_y such that $y \in G_y$ but $x \notin G_y$. It follows that $y \in G_y \subset X - \{x\}$. Hence $X - \{x\}$ is DOS, and to show that $X - \{y\}$ is DOS. Let $x \in X - \{y\}$ this means $x \in X - \{(\phi, \{y\}^c)\} \Rightarrow x \notin X\{y\}^c, x \in X - \{y\}$ and $y \notin X - \{y\}$ then there exists a DOS H_x such that $x \in H_x$ but $y \notin H_x$, it follows that $x \in H_x \subset X - \{y\}$. Hence $X - \{y\}$ is DOS. Accordingly $\{x\}$ is DCS.

4.10. Proposition

For a DTS (X, τ) pairwise T_3 is pairwise T_1 .

Proof : Let DTS (X, τ) be pairwise T_3 , we have $G=(A,B)$, $H=(C,D) \in \tau$ with $x \in G, x \notin H$ and $G \subseteq (\overline{H})$ i.e $A \subseteq D$, take $x \neq y$ in X and $y \in H \Rightarrow y \in D \Rightarrow y \in A \Rightarrow y \in G$ Then (X, τ) is T_1 .

4.11. Proposition

For a DTS (X, τ) pairwise T_3 is pairwise T_2 .

Proof : Let DTS (X, τ) be pairwise T_3 . Take $x \neq y$ in X Since X is pairwise T_1 , then there exist $G \in \tau$ with $x \in G$ and $y \notin G$ and $H \in \tau$ with $y \in H$ and $x \notin H$, and since X is pairwise regular there exist $G, H \in \tau$ such that $x \in G$ and $x \notin H$, $G \subseteq (\overline{H})$ so that $x \in G, y \in H$ and $G \subseteq (\overline{H})$. Accordingly (X, τ) is T_2 .

References

- [1] Atanassov, K, Intuitionistic fuzzy sets, Fuzzy Sets and Systems 20(1),87-96,1986.
- [2] Chang, C. L, Fuzzy Topological Spaces, J. Math. Appl. 24, 182-190, 1968.
- [3] Coker, D, A note on Intuitionistic Sets and Intuitionistic Points, Turkish J. Math. 20(3), 343-351, 1996.

- [4] Coker, D. and Es, A. H. On Fuzzy Compactness In Intuitionistic Fuzzy Topological Spaces, *J. Fuzzy Mathematics* 3(4), 899-909, 1995.
- [5] Coker, D. and Demirci, M. On Intuitionistic Fuzzy Points, *Notes IFS* 1(2), 79-84, 1995.
- [6] P. Fletcher, H. B. Holye and C. W. Patty, The Comparison of Topologies, *Duke Math. J.* 36, 325-331, 1965.
- [7] Garia, J.G. and Rodabaugh, S.E. Order-theoretic, topological, categorical redundancies of interval-valued sets, grey sets, vague sets, interval-valued (intuitionistic) sets, (intuitionistic) fuzzy sets and topologies, *Fuzzy Sets and Systems* 156(3), 445-484, 2005.
- [8] Kelly, J. G, Bitopological Space, *Proc. London Math. Soc.* 13, 71-89, 1963.
- [9] Zadeh, L. A, Fuzzy Sets, *Inform. And Control* 8, 338-353, 1965.