

*Noori Farhan Al-Mayahi                      Duaa Hassan   Hussein*

**University of Al-Qadisiya**  
**College of Computer Science and Mathematics**  
**Department of Mathematics**

### **Abstract**

In this paper we study some new properties of convexly compact sets.

### **Key words**

compact sets and convexly compact sets.

### **1.Introduction**

The genesis of the notation of compactness is connected with the Borel theorem (proved in 1903) stating that every countable open cover of a closed interval admits a finite subcover, and with the Lebesgue observation that the same holds for every open cover of a closed interval ( in [1903] Borel generalization this result, in Lebesgue's setting, to all bounded closed subsets of Euclidean spaces ) [2]. In 2010 Zitkovic [5] introduced the concept of convexly compact sets. A collection  $\Omega$  of sets is said to have the finite intersection property (FIP) [2,4] if the intersection of each finite subcollection of  $\Omega$  is non-empty. A subset  $C$  of a topological space  $X$  is said to be compact if every open cover of  $C$  admits a finite subcover, or equivalently, if and only if every collection of closed subsets of  $C$  with the finite intersection property admits non-empty intersection [2,4]. A function from a topological space into topological space is continuous if and only if the inverse image of every open (closed) set is open (closed) set. A function from a topological space into topological space is said to be closed if the image of every closed set is closed set [2,4].

Let  $\Lambda$  be a non-empty set. The set  $Fin(\Lambda)$  consisting of all non-empty finite subsets of  $\Lambda$  carries a natural structure of a partially ordered set when ordered by inclusion. Moreover, it is a directed set, since  $D_1, D_2 \subseteq D_1 \cup D_2$  for any  $D_1, D_2 \in Fin(\Lambda)$  [5].

**Definition 1.1 [3]**

A topology  $\tau$  on a vector space over a field  $\Phi$  is called a vector topology if the map  $+: X \times X \rightarrow X$  and  $\bullet: \Phi \times X \rightarrow X$  are continuous. A vector space endowed with a vector topology is called a topological vector space.

**Definition 1.2 [1,3]**

A subset  $C$  of a vector space  $X$  is said to be a convex set if  $\lambda x + (1-\lambda)y \in C$  for every  $x, y \in C$  and  $0 \leq \lambda \leq 1$ .

**Remark 1.3 [3]**

- Let  $X$  be a vector space, then
- (i) the empty set and the singleton set are convex sets.
  - (ii) every intersection of convex sets is a convex set.
  - (iii) The closure of every convex set is a convex set.

**Theorem 1.4 [1,3]**

- (i) The image of every convex set under a linear map is convex.
- (ii) The inverse image of a convex set under a linear map is convex.

**Definition 1.5 [2,4]**

Let  $X$  be any non-empty set. A filter on  $X$  is a non-empty collection  $\mathbf{F}$  of subsets of  $X$  satisfying the following axioms.

- [F1]  $\emptyset \notin \mathbf{F}$ .
- [F2] If  $F \in \mathbf{F}$  and  $H \supset F$ , then  $H \in \mathbf{F}$ .
- [F3] If  $F \in \mathbf{F}$  and  $H \in \mathbf{F}$ , then  $F \cap H \in \mathbf{F}$ .

**Remark 1.6 [2,4]**

Every filter on a non-empty set  $X$  admits the finite intersection property.

**Definition 1.7 [2,4]**

Let  $(X, \tau)$  be a topological space and let  $\mathbf{F}$  be a filter on  $X$ . The point  $x \in X$  is said to be a cluster point of  $\mathbf{F}$  if and only if  $x \in \overline{F}$  for all  $F \in \mathbf{F}$ .

**2. Convexly compact sets**

In this section we shall study some new properties of convexly compact sets.

**Definition 2.1 [ 5]**

A convex subset  $C$  of a topological vector space  $X$  is said to be convexly compact if for any non-empty set  $A$  and any family  $\{F_\alpha : \alpha \in A\}$  of closed and convex subsets of  $C$ , the condition

$$\forall D \in \text{Fin}(A), \bigcap_{\alpha \in D} F_\alpha \neq \phi \tag{2.1}$$

implies

$$\bigcap_{\alpha \in A} F_\alpha \neq \phi, \tag{2.2}$$

Without the additional restriction that the sets  $\{F_\alpha : \alpha \in A\}$  be convex, Definition 2.1—postulating the finite-intersection property for families of closed and convex sets would be equivalent to the classical definition of compactness. It is, therefore, immediately clear that any convex and compact subset of a topological vector space is convexly compact [5]. The converse however is not true ( see [5] Example 2.2).

**Definition 2.2.**

A topological vector space  $X$  is said to be convexly compact if for any non-empty set  $A$  and any family  $\{F_\alpha\}_{\alpha \in A}$  of closed and convex subsets of  $X$ , the condition

$$\forall D \in \text{Fin}(A), \bigcap_{\alpha \in D} F_\alpha \neq \phi \tag{2.1)*}$$

implies

$$\bigcap_{\alpha \in A} F_\alpha \neq \phi, \tag{2.2)*}$$

**Definition 2.3**

A subset of a vector space  $X$  is said to be co-convex if its complement is convex.

**Theorem 2.4**

If  $A$  is co-convex subset of a vector space  $X$ , then  $\sigma A$  is co-convex for every  $\sigma \in \Phi / \{0\}$ .

**Proof**

we need to prove that if  $X/A$  is convex set, then  $X/\sigma A$  is convex set. Let  $x, y \in X/\sigma A$  and  $0 \leq \lambda \leq 1$ . Then

$$\begin{aligned} x, y \in X \text{ and } x, y \notin \sigma A \\ \Rightarrow x, y \in X \text{ and } \sigma^{-1}x, \sigma^{-1}y \notin A. \end{aligned}$$

$\Rightarrow \sigma^{-1}x, \sigma^{-1}y \in X$  ( since  $X$  is a vector space and  $\sigma^{-1} \in \Phi$ ) and  $\sigma^{-1}x, \sigma^{-1}y \notin A$ . Therefore  $\sigma^{-1}x, \sigma^{-1}y \in X/A$ . Since  $X/A$  is convex set then

$$\begin{aligned} \lambda(\sigma^{-1}x) + (1-\lambda)(\sigma^{-1}y) &\in X/A \\ \Rightarrow \sigma^{-1}(\lambda x) + \sigma^{-1}((1-\lambda)y) &\in X/A \\ \Rightarrow \sigma^{-1}(\lambda x + (1-\lambda)y) &\in X/A \\ \Rightarrow \sigma^{-1}(\lambda x + (1-\lambda)y) \in X \text{ and } \sigma^{-1}(\lambda x + (1-\lambda)y) &\notin A \\ \Rightarrow \lambda x + (1-\lambda)y \in X \text{ and } \lambda x + (1-\lambda)y &\notin \sigma A \\ \Rightarrow \lambda x + (1-\lambda)y \in X/\sigma A. \end{aligned}$$

Then  $X/\sigma A$  is convex and this complete the proof.

**Remark 2.5**

The above theorem dose not still true if  $\sigma = 0$ . For example, consider the real line and let  $A$  be any co-convex subset of the real line. Since  $\sigma A = \{0\}$  then  $(\sigma A)^c = (-\infty, 0) \cup (0, \infty)$  is not convex . Indeed the line segment that joint  $-1$  and  $1$  ( for example ) dose not lie in  $(\sigma A)^c$ .

**Theorem 2.6**

The arbitrary union of a collection of a co-convex set is co-convex.

**Proof**

Suppose that  $\{A_\sigma : \sigma \in \Lambda\}$  be arbitrary collection of a co-convex subsets of a vector space  $X$ . Then The collection  $\{A_\sigma^c : \sigma \in \Lambda\}$  is convex for every  $\sigma \in \Lambda$ . By Remark 1.3 (ii) we get  $\bigcap_{\sigma \in \Lambda} A_\sigma^c$  is convex set. By De-Morgan Law we have

$$\bigcap_{\sigma \in \Lambda} A_\sigma^c = \left( \bigcup_{\sigma \in \Lambda} A_\sigma \right)^c .$$

Therefore  $\left( \bigcup_{\sigma \in \Lambda} A_\sigma \right)^c$  is convex i.e.  $\bigcup_{\sigma \in \Lambda} A_\sigma$  is co-convex and the proof is complete.

**Theorem 2.7**

A topological vector space  $X$  is convexly compact if and only if every co-convex open cover of  $X$  admits a finite co-convex open cover.

**Proof**

Suppose that  $X$  is convexly compact and let  $\{G_\alpha : \alpha \in \Lambda\}$  be a co-convex open cover of  $X$ , so that

$$X = \bigcup_{\alpha \in \Lambda} G_\alpha .$$

Then

$$\phi = \bigcap_{\alpha \in \Lambda} G_\alpha^c$$

Thus  $\{G_\alpha^c : \alpha \in \Lambda\}$  be a collection of convex closed sets with empty intersection so by hypothesis there exists  $D \in Fin(\Lambda)$  such that  $\bigcap_{\alpha \in D} G_\alpha^c = \phi$ . Thus

$$\left( \bigcup_{\alpha \in D} G_\alpha \right)^c = \phi \Rightarrow \bigcup_{\alpha \in D} G_\alpha = X .$$

Conversely, suppose that for every co-convex open cover of  $X$  admits a finite co-convex open cover, and let  $\{F_\alpha : \alpha \in \Lambda\}$  be a collection of convex closed subset of  $X$  such that condition (2.1)\* holds, i.e.

$$\forall D \in Fin(A), \bigcap_{\alpha \in D} F_\alpha \neq \phi .$$

Suppose if possible,  $\bigcap_{\alpha \in A} F_\alpha = \phi$ . Then

$$X = \left( \bigcap_{\alpha \in \Lambda} F_\alpha \right)^c = \bigcup_{\alpha \in \Lambda} F_\alpha^c .$$

This means that  $\{F_\alpha^c : \alpha \in \Lambda\}$  is a co-convex open cover of  $X$ . By hypothesis, there exists  $D \in Fin(\Lambda)$  such that

$$\bigcup_{\alpha \in D} F_{\alpha}^c = X \Rightarrow \left( \bigcap_{\alpha \in D} F_{\alpha} \right)^c = X \Rightarrow \bigcap_{\alpha \in D} F_{\alpha} = \phi$$

But this contradicts the condition (2.1)\*. Hence we must have  $\bigcap_{\alpha \in \Lambda} F_{\alpha} \neq \phi$ .

### Theorem 2.8

A topological vector space  $X$  is convexly compact if and only if any basic co-convex open cover of  $X$  admits a finite subcover.

#### Proof

Let  $X$  be a convexly compact space. Then by Theorem 2.7 every co-convex open cover of  $X$  admits a finite subcover. In particular, every basic co-convex open cover of  $X$  admit a finite subcover.

Conversely, suppose that every basic co-convex open cover of  $X$  admits a finite subcover. Let  $\{C_{\lambda} : \lambda \in \Lambda\}$  be any co-convex open cover of  $X$ . If  $\{B_{\gamma} : \gamma \in \Gamma\}$  be any co-convex open base for  $X$ , then each  $C_{\lambda}$  is a union of some members of  $B$ .

That is there exists  $\Gamma_{\lambda} \subseteq \Gamma$  such that  $C_{\lambda} = \bigcup_{\gamma \in \Gamma_{\lambda}} B_{\gamma}$  for every  $\lambda \in \Lambda$ . (Note that by

Theorem 2.6 we have the union of co-convex sets is also co-convex set). And the totality of all such members of  $B$  is evidently a basic co-convex open cover of  $X$ . By hypothesis this collection of members of  $B$  admits a finite subcover, say,  $\{B_{\gamma_i} : i=1,2,\dots,n\}$ . For each  $B_{\gamma_i}$  in this finite subcover, we can select a  $C_{\lambda_i}$  from  $\Omega$  such that  $B_{\gamma_i} \subseteq C_{\lambda_i}$ . It follows that the finite subcollection  $\{C_{\lambda_i} : i=1,2,\dots,n\}$  each arise in this way is a subcover of  $\Omega$ . Hence  $X$  is convexly compact.

### Theorem 2.9

If  $X$  be a convexly compact topological vector space. Then every convex filter on  $X$  admit a cluster point.

#### Proof

Assume  $X$  be a convexly compact topological vector space and let  $\Phi$  be any convex filter on  $X$ . Then  $\Phi$  be a collection of convex subsets of  $X$  and by Remark 1.6 it has FIP. Since by Remark 1.3(iii) the closure of convex set is also convex and the closure of any set is closed, then  $\{\overline{F} : F \in \Phi\}$  is a collection of convex closed subsets of  $X$ . Since  $\Phi$  has FIP property, then so is  $\{\overline{F} : F \in \Phi\}$ .

By hypothesis  $\bigcap_{F \in \Phi} \overline{F} \neq \emptyset$ , hence there exists at least one point  $p \in \bigcap_{F \in \Phi} \overline{F}$ . This implies that  $p \in \overline{F}$  for every  $F \in \Phi$ . Hence  $p$  is a cluster point of  $\Phi$  by Definition 1.7.

### Theorem 2.10

The image of convexly compact set under a continuous bijective linear map is convexly compact.

#### Proof

Suppose that  $C$  be a convexly compact in a topological vector space  $X$  and let  $f$  be continuous linear map from  $X$  onto another topological vector space  $Y$ . Since  $C$  is convex in  $X$  and  $f$  is linear then by Theorem 1.4 (i)  $f(C)$  is convex set in  $Y$ . To show that  $f(C)$  is a convexly compact. Let  $\{F_\alpha : \alpha \in \Lambda\}$  be a collection of closed and convex subsets of  $f(C)$  with the property that

$$\bigcap_{\alpha \in D} F_\alpha \neq \emptyset, \forall D \in \text{Fin}(\Lambda).$$

Since  $f$  is continuous, and  $F_\alpha$  is closed  $\forall \alpha \in \Lambda$  then  $\{f^{-1}(F_\alpha) : \alpha \in \Lambda\}$  is closed. Since  $f$  is linear, and  $F_\alpha$  is convex  $\forall \alpha \in \Lambda$  then by theorem 1.4(ii)  $\{f^{-1}(F_\alpha) : \alpha \in \Lambda\}$  is convex. Since  $F_\alpha \subset f(C)$  for every  $\alpha$ , then  $f^{-1}(F_\alpha) \subset C$  for every  $\alpha$ . Hence  $\{f^{-1}(F_\alpha) : \alpha \in \Lambda\}$  is collection of convex and closed subsets of  $C$  since  $\bigcap_{\alpha \in D} F_\alpha \neq \emptyset$  and  $f$  is onto then  $f^{-1}(\bigcap_{\alpha \in D} F_\alpha) \neq \emptyset \forall \alpha \in D$ . Since  $f$  is bijective then  $f^{-1}(\bigcap_{\alpha \in D} F_\alpha) = \bigcap_{\alpha \in D} f^{-1}(F_\alpha)$  and hence  $\bigcap_{\alpha \in D} f^{-1}(F_\alpha) \neq \emptyset \forall D \in \text{Fin}(A)$ . Since  $C$  is convexly compact, then  $\bigcap_{\alpha \in A} f^{-1}(F_\alpha) \neq \emptyset$  then  $f(\bigcap_{\alpha \in A} f^{-1}(F_\alpha)) \neq \emptyset$  since  $f(\bigcap_{\alpha \in A} f^{-1}(F_\alpha)) = \bigcap_{\alpha \in A} f(f^{-1}(F_\alpha)) = \bigcap_{\alpha \in A} F_\alpha$  [since  $f$  is bijective]. Thus  $\bigcap_{\alpha \in A} F_\alpha \neq \emptyset$ .  $\Rightarrow f(C)$  is convexly compact. This complete the proof.

### Theorem 2.11

The inverse image of every convexly compact set under closed bijective linear map is convexly compact.

**Proof**

Suppose that  $X$  and  $Y$  are two topological vector spaces and  $f : X \rightarrow Y$  be a closed bijective linear map. Let  $C$  be a convexly compact set in  $Y$ . To show that  $f^{-1}(C)$  is convexly compact in  $X$ . Since  $C$  is convex and  $f$  is linear then by Theorem 1.4 (ii)  $f^{-1}(C)$  is convex set. Now, let  $\{F_\alpha : \alpha \in \Lambda\}$  be a collection of convex closed subsets of  $f^{-1}(C)$  with the property that  $\bigcap_{\alpha \in D} F_\alpha \neq \phi$ , for every  $D \in \text{Fin}(\Lambda)$ . Since  $F_\alpha \subseteq f^{-1}(C)$  and  $f$  is onto, then  $f(F_\alpha) \subseteq f(f^{-1}(C)) = C$ . Since  $F_\alpha$  is closed  $\forall \alpha \in \Lambda$  and  $f$  is closed function, then  $f(F_\alpha)$  is closed subset of  $C$ . Since  $F_\alpha$  is convex  $\forall \alpha \in \Lambda$  and  $f$  is linear then  $f(F_\alpha)$  is convex set. Thus  $\{f(F_\alpha) : \alpha \in \Lambda\}$  is a collection of closed convex subsets of  $C$ . Since  $\bigcap_{\alpha \in D} F_\alpha \neq \phi, \forall D \in \text{Fin}(\Lambda)$  then  $f(\bigcap_{\alpha \in D} F_\alpha) \neq \phi$ . But  $f$  is onto, thus  $f(\bigcap_{\alpha \in D} F_\alpha) \subseteq \bigcap_{\alpha \in D} f(F_\alpha)$ , therefore  $\bigcap_{\alpha \in D} f(F_\alpha) \neq \phi$  i.e.  $\{f(F_\alpha) : \alpha \in \Lambda\}$  admits finite intersection property. By convexly compactness of  $C$ , we have  $\bigcap_{\alpha \in \Lambda} f(F_\alpha) \neq \phi$ . Since  $f$  is onto, then  $f^{-1}(\bigcap_{\alpha \in \Lambda} f(F_\alpha)) \neq \phi$  since  $f$  is one-one then  $f^{-1}(\bigcap_{\alpha \in \Lambda} f(F_\alpha)) = \bigcap_{\alpha \in \Lambda} f^{-1}(f(F_\alpha)) = \bigcap_{\alpha \in \Lambda} F_\alpha$ ; hence  $\bigcap_{\alpha \in \Lambda} F_\alpha \neq \phi$ , i.e.  $f^{-1}(C)$  is convexly compact, and this complete the proof.

**Theorem 2.12**

If  $A$  is convexly compact subset of a topological vector space  $X$ . Then the set  $\lambda A$  is convexly compact where  $\lambda \in \Phi$ .

**Proof**

Suppose that  $A$  is convexly compact subset of a topological vector space  $X$  and  $\lambda \in \Phi$ . Since the function  $\bullet : \Phi \times X \rightarrow X$  which define by  $(\lambda, x) = \lambda x$  is continuous by definition of a topological vector space, then by Theorem 2.10 the set  $\lambda A$  is convexly compact and the proof is complete.

## **References**

- [1] G. Zitkovic, " Convexly Compactness and its Applications" Math. Finan Econ 3(2010). 1-12.
- [2] J.N.Sharma,"Topology" Published by Krishna Pracushna , Mandir , and printed at Mano, 1977.
- [3] N.F. Al-Mayahi and A.H. Battor " *Introduction to Functional Analysis*" Iraq. Najef. 2005.
- [4] R. Engelking " General Topology " polish. Scientific publishers. Warsawx 1989.
- [5] W .Rudin ,"*Functional Analysis*", New York. 1973