

Korovkin Type Theorem in Weighted Spaces ($L_{p,\alpha}$ -Space)

Alaa H. Al-Abdulla

**Ministry of Education, Baghdad Education Directorate, Al-Karkh
the Second, Department of Specialized Supervision**

Recived :15\7\2014

Revised : 13\5\2015

Accepted :25\8\2015

Abstract:

In this study, we obtain some Korovkin type approximation theorem by positive linear operators on $L_{p,\alpha}$ -space of unbounded functions by using the test functions $\{1,x,x^2\}$.

Mathematics Subject Classification: 46S40 .

Introduction:

The approximation problem arises in many contexts of “Numerical Analysis and Computing”. The “Quadrature” is one such context, for example. Wierstrass (1885) proved his celebrated approximation theorem: If $f \in C[a,b]$; for $\delta > 0$ there is a polynomial p such that $|f - p| < \delta$. In other words, the result established the existence of an algebraic polynomial in the relevant variable.

The great Russian mathematician S. N. Bernstein proved the Weirstrass theorem in such a manner as was very stimulating and interesting in many ways.

As we know, approximation theory has important applications in various areas of mathematics (see [3,4,7]). Most of the classical operators tend to converge to the value of the function being approximated.

It is also known J. P. King [6] has presented an unexpected example of operators of Bernstein type which preserve that test function e_0 and e_2 of Bohman-Korovkin theorem.

Now, we constructive approximation of unbounded functions over L_n which is a linear positive operators from $B_\alpha = \{f : A \longrightarrow \mathbb{R} \text{ such that } |f(x)e^{-\alpha x}| \leq M\}$

into itself, we use the special norm:

$$\|f\|_{p,\alpha} = \left(\int_A |f(x)e^{-\alpha x}|^p dx \right)^{1/p} < \infty, \alpha > 0$$

Here, we define the weighting modulus of $f \in B_\alpha$ by:

$$\omega(f, \delta)_{p,\alpha} = \left(\int_A \left| \frac{1}{h} \Delta f(x) e^{-\alpha x} \right|^p dx \right)^{1/p}$$

where $\frac{1}{h} \Delta f(x) = |f(x) - f(y)|$, $|x - y| < h$ for all $h < \delta$.

Main Results:

By using Korovkin theorem in [7],[5],[3], we want to estimate the degree of best approximation of unbounded function in terms of weighted modulus of function.

Theorem:

Let f be unbounded real-valued function on $[a,b]$, let $L_n : B_\alpha \longrightarrow B_\alpha$ be a linear positive operator, such that:

$$\|L_n(g) - g\|_{p,\alpha} \longrightarrow 0, \text{ for } n \longrightarrow \infty \text{ and } g \in \{1, x, x^2\}$$

Then:

$$\|L_n(f) - f\|_{p,\alpha} \longrightarrow 0, \text{ for } n \longrightarrow \infty, \text{ for each } f \in B_\alpha$$

Proof:

From [5] , $\|L_n(g) - g\|_{p,\alpha} \longrightarrow 0$ for $n \longrightarrow \infty$, we get:

$$L_n(g(y))(x) = g(x) + \omega\left((g(y))(x), \delta \right)_{p,\alpha},$$

where $\omega\left((g(y))(x), \delta \right)$ represents the error we make in approximating $g(x)$ with $L_n(g(y))(x)$, and hence it is such that:

$$\|\omega((g(y))(x), \delta)\|_{p,\alpha} \longrightarrow 0$$

Now, we have to prove that:

$$\|L_n(f) - f\|_{p,\alpha} \longrightarrow 0, \text{ for } n \longrightarrow \infty, \forall f \in B_\alpha$$

we fixed $f \in B_\alpha$ and $\omega(f, \delta)_{p,\alpha} > 0$, we want to prove that there exist \bar{n} (as a function of f and ω), such that for any $n \geq \bar{n}$ and for every $x \in \mathbb{R}$, we have:

$$|(L_n(f) - f)e^{-\alpha x}| < \omega(f, \delta)_{p,\alpha}.$$

And we write:

$$\begin{aligned} \Delta_n(f)(x) &= |(L_n(f(y))(x) - f(x))e^{-\alpha x}| \\ &= |(L_n(f(y))(x) - 1.f(x))e^{-\alpha x}| \\ &= |(L_n(f(y))(x) - [L_n(1)(x) - \omega(f, \delta)_{p,\alpha}(x)]f(x))e^{-\alpha x}| \end{aligned}$$

since $1 = L_n(1)(x) - \omega(f, \delta)_{p,\alpha}(x)$

$$\begin{aligned} \Delta_n(f)(x) &\leq |(L_n(f(y))(x) - L_n(1)f(x))e^{-\alpha x}| + |f(x)|\omega(f, \delta)_{p,\alpha}e^{-\alpha x} \\ &\leq |(L_n(f(y))(x) - L_n(1)f(x))e^{-\alpha x}| + |f(x)|e^{-\alpha x} \|\omega(f, \delta)_{p,\alpha}\| \end{aligned}$$

$$\leq |(L_n(f(y))(x) - L_n(1)f(x))e^{-\alpha x}| + \|f(x)\| \|\omega(f, \delta)_{p,\alpha}\| \text{ and from [5] and [2]}$$

we have

$$\leq |(L_n(f(y)))(x) - L_n(1)f(x)e^{-\alpha x}|_+ \left(\int_A |f(x)e^{-\alpha x}|^p M\omega(f, \delta)_\alpha(x) dx \right)^{1/p}$$

$$= |(L_n(f(y)))(x) - L_n(1)f(x)e^{-\alpha x}|_+ \|f\|_{p,\alpha} \|w(f, \delta)_\alpha\|_p$$

we have to find:

$$\Delta_n(f)(x) \leq L_n \left(|f(y) - f(x)| e^{-\alpha x} \right) (x) |f(y) - f(x)| e^{-\alpha x} + \frac{\bar{\omega}}{4}, \forall n \geq \bar{n}$$

Whenever $\|w(f, \delta)_\alpha\|_p \rightarrow 0$ for $n \rightarrow \infty$. Now:

$$\left(\int_A |f(y)e^{-\alpha y} - f(x)e^{-\alpha x}|^p dx \right)^{1/p} = \|\Delta_n f\|_{p,\alpha}$$

$$= \omega(f, \delta)_{p,\alpha}, \text{ where } \delta = |x - y|$$

and $\omega(f, \delta)_{p,\alpha} \rightarrow 0$. If $|x - y| \rightarrow 0$, then from [1] we have:

$$|f(x)e^{-\alpha x} - f(y)e^{-\alpha y}| < \bar{\omega} \chi\{z : |z - x| < \delta\}(y) + 2\|f\|_{p,\alpha} \chi\{z : |z - x| \geq \delta\}(y),$$

$$\text{Now, } |z - x| \geq \delta \Leftrightarrow \frac{|z - x|}{\delta} \geq 1 \Rightarrow \frac{(z - x)^2}{\delta^2} \geq 1$$

In conclusion, in the set the inequality is satisfied, we can write:

$$\chi\{x : |z - x| \geq \delta\}(y) = 1 \leq \frac{(y - x)^2}{\delta^2}$$

otherwise, we have:

$$\chi\{x : |z - x| \geq \delta\}(y) = 0 \leq \frac{(y - x)^2}{\delta^2}$$

Therefore, uniformly with respect to $x, y \in \mathbb{R}$ for any $\bar{\omega} > 0$, we find $\delta = \delta \bar{\omega}$, such that:

$$|f(x)e^{-\alpha x} - f(y)e^{-\alpha y}| \leq \bar{\omega} (1) + 2\|f\|_{p,\alpha} \frac{(y - x)^2}{\delta^2}$$

Choose $\bar{\omega} = \frac{\omega}{4}$.

By applying the operator $L_n(\cdot)$ for every $n \geq \bar{n}_1$, we find that:

$$\begin{aligned} \Delta_n(f)(x) &\leq \frac{\omega}{4} + L_n \left(\bar{\omega} + \frac{2 \|f\|_{p,\alpha}}{\delta^2} (y-x)^2 \right) (x) \\ &= \frac{\omega}{4} + \bar{\omega} L_n(1) + \frac{2 \|f\|_{p,\alpha}}{\delta^2} L_n \left(y^2 - 2yx + x^2 \right) (x) \end{aligned}$$

Hence, $L_n(1)(x) = \left(1 + \omega(1, \delta)_{p,\alpha} \right) (x)$ and the constant 1 is a test function.

Therefore, from [3], [5] we have:

$$\begin{aligned} \Delta_n(f)(x) &\leq \frac{\omega}{4} + \bar{\omega} \left(1 + \omega(1, \delta)_{p,\alpha}(x) \right) + \frac{2 \|f\|_{p,\alpha}}{\delta^2} \left\{ x^2 + \omega_\alpha(y^2, \delta)(x) - 2x \left[x + \right. \right. \\ &\quad \left. \left. \omega_\alpha(y, \delta)(x) \right] + x^2 \left(1 + \omega_\alpha(1, \delta)(x) \right) \right\} \end{aligned}$$

Now, by definition of limit, there exist a functions \bar{n}_2, \bar{n}_3 such that for $n \geq \bar{n}_2$, we have

$\omega_\alpha(1, \delta) \leq 1$, uniformly with respect to $x \in \mathbb{R}^n$ and for $n \geq \bar{n}_3$, we have:

$$\|f\|_{p,\alpha} \|\omega_\alpha(y^2, \delta)(x) - 2x\omega_\alpha(y, \delta)(x) + x^2\omega_\alpha(1, \delta)(x)\|_{p,\alpha} \leq \frac{\delta^2 \bar{\omega}}{2}$$

Now, by taking $n \geq \bar{n}$, with $\bar{n} = \max\{\bar{n}_1, \bar{n}_2, \bar{n}_3\}$ (depending on both f and ω), we have:

$$\begin{aligned} \Delta_n(f)(x) &\leq \frac{\omega}{4} + 2\bar{\omega} + \frac{2 \|f\|_{p,\alpha}}{\delta^2} \{ \omega_\alpha(y^2, \delta)(x) - 2x\omega_\alpha(y, \delta)(x) + x^2\omega_\alpha(1, \delta)(x) \} \\ &\leq \frac{\omega}{4} + 3\bar{\omega} = 1 \end{aligned}$$

References:

- [1] O. Agratini and O. Doğru, “Weighted Approximation by q-Szasz-King Type Operators”, Vol.14, No.4, pp.1283-1296, August, 2010.
- [2] F. Cao and Y. Lin, “Approximation Theorems by Positive Linear Operators in Weighted Spaces”, Positivity, 15: 87-103 (2011).
- [3] F. Itomre, M. Cmpiti, “Korovkin Type Approximation Theory and its Application”, Walter de Gryter, Berlin (1994).
- [4] R. A. Devore, The Approximation of Continuous Functions by Positive Linear Operators”, Lecture Notes in Mathematics, Vol.293, Springer, Berlin (1972).
- [5] F. Didone, M. Giannuzzi and L. Paulon, “Korovkin Theorems and Applications in Approximation Theory and Numerical Algebra”, March 14, 2011.
- [6] J. P. King, “Positive Liner Operators which Preserve x^2 ”, Acta Math. Hungar, 99(3) (2003), 208.
- [7] P. P. Korovkin, “Liner Operators and Approximation Theory”, Hindustan Public Corporation, Delhi (1960).
- [8] S. A. Wahid, “Efficient Approximation Using Probabilistically Improved Combination Structure of Bernstein’s Polynomial Operators Weights Through the Fusion of Dual-Perspectives”, Vol.3, No.3, August (2011).

نوع معين من نظرية كورفكين في الفضاء الوزني ($L_{p,\alpha}$ -Space)

الآء حميد علون العبدالله

وزارة التربية العراقية، تربية بغداد الكرخ الثانية، قسم الإشراف الإختصاصي

المستخلص:

في هذه الدراسة إستنتجنا نوع من تقريب كورفكين بواسطة المؤثرات الخطية الموجبة للدوال غير المقيدة في الفضاء الوزني ($L_{p,\alpha}$ -Space).